

Consistent Application of Maximum Entropy to Quantum-Monte-Carlo Data

W. von der Linden^a, R. Preuss^b, and W. Hanke^b

a) Max-Planck-Institut für Plasmaphysik, EURATOM Association

D-85740 Garching b. München, Germany

e-mail: wvl@ibmop5.ipp-garching.mpg.de

b) Physikalisches Institut, Universität Würzburg, Am Hubland

D-97074 Würzburg, Federal Republic of Germany

e-mail: roland@physik.uni-wuerzburg.de

(February 1, 2008)

Abstract

Bayesian statistics in the frame of the maximum entropy concept has widely been used for inferential problems, particularly, to infer dynamic properties of strongly correlated fermion systems from Quantum-Monte-Carlo (QMC) imaginary time data. In current applications, however, a consistent treatment of the error-covariance of the QMC data is missing. Here we present a closed Bayesian approach to account consistently for the QMC-data.

PACS numbers: 71.20.Ad

I. INTRODUCTION

Bayesian statistics [1] provides a general and consistent frame for logical inference based on incomplete and noisy data and prior knowledge. Combined with the entropic *Prior* it is referred to as quantified maximum entropy [2,3] (MaxEnt) and yields the most probable and unbiased result, given the data-constraints and prior knowledge. MaxEnt has originally been introduced [2,3] to infer celestial images from incomplete and noisy radio-astronomic data. In the sequel, it has been applied successfully to various other data-analysis problems [4–6,8].

Here we will focus on the ill-posed inversion problem encountered in Quantum-Monte-Carlo (QMC) simulations [5,9]. In this field, MaxEnt has become a standard and successful technique to infer dynamic properties of strongly correlated fermion systems from imaginary time QMC data, which intrinsically is an inverse Laplace transform. QMC yields values for dynamic quantities along the imaginary time-axis for a finite number of times. The inversion is not unique due to the limited number of data and the presence of statistical errors. A direct inversion of the Laplace transform would tremendously overfit the noise and the desired signal would be buried underneath it. Bayesian probability theory provides a consistent frame to separate the signal from the noise.

A complication arises, however, in the present application as the errors of the QMC-data are correlated. It has been proposed [5] to include the error-covariance matrix as exact data-constraints. This approach has been heavily debated as it leads to the following dilemma. For a standard QMC sample size, the off-diagonal elements of the covariance matrix are not negligible and ought to be taken into account. However, the errors of the covariance matrix are huge and the information provided by the QMC covariance matrix is useless and in many applications even disadvantageous. The error of the covariance matrix can be decreased by increasing the sample size. This procedure is, however, computationally very expensive and, moreover, needless as the correlation of the errors decreases at the same time. The dilemma obviously arises due to the neglect of the errors of the covariance matrix. Here we present a

closed Bayesian approach to account fully for the noisy QMC data plus covariance matrix. This scheme allows to infer reliable results with the least amount of computer time.

Bayes' theorem aims at determining the posterior probability (*posterior*) $P(A|DCH)$ of the sought-for quantity A , given hypotheses H and QMC data D , which are related to A via

$$D_l^{ex} = D_l^{ex}(\{A\}) \quad . \quad (1.1)$$

The experimental data D_l deviate from the exact values by the statistical QMC errors η_l . Due to the QMC-algorithm, the errors are correlated and the information about the error-covariance matrix will be denoted by C . Bayes' theorem relates the *posterior* to the likelihood function $P(D|ACH)$, which contains the error statistic of the QMC data, and the prior probability $P(A|CH)$

$$P(A|DCH) = \frac{P(D|ACH)P(A|CH)}{P(D|CH)} \quad . \quad (1.2)$$

The most honest *Prior* should summarize all our prior knowledge – the knowledge we have about A prior to receiving the experimental data A – and nothing more, i.e. it should be as ignorant as possible otherwise [1]. Our prior knowledge is part of the hypotheses H . If nothing is known about the solution A the most ignorant *Prior* would simply be $P(A|CH) = \text{const.}$ (If nothing is known about a coin which we are going to toss, common sense tells us to assign equal prior probabilities to the events "head (tail) shows up").

In the case of a positive, additive distribution function (PAD) – e.g. the spectral density – the most ignorant *Prior* is the entropic *Prior* [10]

$$P(A|CH) = \frac{1}{Z_S} \exp\left(\alpha \underbrace{\int [A(\omega) - m(\omega) - A(\omega) \ln(\frac{A(\omega)}{m(\omega)})] d\omega}_S\right) \quad . \quad (1.3)$$

Z_S is the normalization constant guarantying $\int P(A|CH) \mathcal{D}A = 1$. In the following we will assume, as indicated in Eq. 1.3 that $A = A(\omega)$ is a function of the frequency ω . The entropy S is measured relative to a default-model $m(\omega)$ which contains "weak prior assumptions" which can still be overruled by the data constraints.

Next we will turn to the determination of the *Likelihood*, the central topic of this paper. Usually this part is dealt with in one sentence: “the *Likelihood* function is given by a Gaussian”. Here we will be more specific why and when this is true. The *Likelihood* quantifies the probability for the realization of the specific data-values D measured in the experiment, supposing the exact function A were known. Given A , the exact values D^{ex} in data-space are also known (Eq. 1.1). The *Likelihood* describes therefore the error statistics of the QMC-data

$$P(D|ACH) =: \rho(D^{\text{ex}} - D) = \rho(\eta) \quad \text{given C and H} \quad . \quad (1.4)$$

In the following, C stands for the error-covariance matrix C_{ij}

$$C_{ij} = \frac{1}{N-1} \langle \eta_i \eta_j \rangle = \frac{1}{N-1} \langle (D_i^{\text{ex}} - D_i)(D_j^{\text{ex}} - D_j) \rangle \quad , \quad (1.5)$$

measured by QMC. N is the number of data-values D_i .

To begin with, we assume that the exact values of the error-covariance are known. The resulting problem is to determine the PAD $\rho(\eta)$ given the constraints

$$C_{ij} = \int \rho(\eta) \eta_i \eta_j d\eta \quad (1.6)$$

$$1 = \int \rho(\eta) d\eta \quad . \quad (1.7)$$

This is a problem falling into the realm of Jaynes’ MaxEnt [11], which is analogous to deriving the barometric formula, Maxwell’s velocity distribution, or Fermi- and Bose statistics. In this framework, $\rho(\eta)$ is obtained upon maximizing the entropy, subject to the exact data-constraints. Treating the constraints with Lagrange parameters we are out for the maximum of

$$\mathcal{L} = S - \sum_{ij} \lambda_{ij} \left(\int \rho(\eta) \eta_i \eta_j d\eta - C_{ij} \right) - \lambda_0 \left(\int \rho(\eta) d\eta - 1 \right) \quad , \quad (1.8)$$

with λ_0, λ_{ij} being Lagrange parameters. Upon maximizing $\mathcal{L}$ with respect to $\rho(\eta)$ one obtains an analytic expression for the solution

$$\rho(\eta) = \frac{1}{Z} e^{-\sum_{ij} \lambda_{ij} \eta_i \eta_j} \quad . \quad (1.9)$$

An ignorant, flat default model ($\int m(\omega)d\omega = 1$) has been assumed. Z is determined via the normalization constraint

$$Z = \int e^{-\sum_{ij} \lambda_{ij} \eta_i \eta_j} d^N \eta = \frac{\pi^{N/2}}{\sqrt{\det(\lambda_{ij})}} \quad (1.10)$$

The covariance constraint implies

$$C_{ij} = -\frac{\partial \ln(Z)}{\partial \lambda_{ij}} = \frac{1}{2}(\lambda^{-1})_{ij} \quad \Rightarrow \quad \lambda_{ij} = \frac{1}{2}(C^{-1})_{ij} \quad . \quad (1.11)$$

Hence the *Likelihood* is the ubiquitous normal distribution

$$\rho(\eta) = \frac{1}{\sqrt{\det(2\pi C)}} e^{-\frac{1}{2} \sum_{ij} \eta_i C_{ij}^{-1} \eta_j} \quad (1.12)$$

which simplifies to a Gaussian if the errors are uncorrelated $C_{ij} = \delta_{ij} \sigma_i^2$

$$\rho(\eta) = \frac{1}{\sqrt{\prod (2\pi \sigma_i^2)}} e^{-\frac{1}{2} \sum_i \frac{\eta_i^2}{\sigma_i^2}} \quad (1.13)$$

Unfortunately, this handy result is only valid if the QMC error-covariance were known exactly, which is not the case. Therefore, the errors of the covariance matrix have to be treated on the same footing as the errors of the QMC data D in the first place – by quantified MaxEnt [2].

Again, the *posterior* for $\rho(\eta)$, given the QMC error-covariance C_{ij} , and the statistical errors σ_{ij} of C_{ij} and all our hypotheses can be determined via Bayes' theorem

$$P(\rho|C\sigma H) = \frac{P(C|\rho\sigma H)P(\rho|H)}{P(C|H)} \quad . \quad (1.14)$$

Superfluous conditions have been discarded. Again the entropic *Prior* is invoked. QMC simulations provide the statistical error σ_{ij} of the covariance matrix. Further information is not available by present QMC simulations. We therefore assume that the error σ_{ij} are uncorrelated and known, this is part of our hypotheses H . A generalization beyond this assumption is straightforward. Along the lines presented above the *Likelihood* reads

$$P(C|\rho\sigma H) = \exp\left(-\frac{1}{2} \underbrace{\sum_{ij} \frac{(C_{ij} - \int \rho(\eta) \eta_i \eta_j d\eta)^2}{\sigma_{ij}^2}}_{\chi^2}\right) \quad . \quad (1.15)$$

The MaxEnt result is hence obtained upon maximizing the *posterior* $P(\rho|C\sigma H) \propto \exp(\alpha S - \frac{1}{2}\chi^2)$ or rather

$$\mathcal{L}(\rho, C) = \alpha S - \frac{1}{2}\chi^2 \quad (1.16)$$

The solution can be cast into the same functional form (Eq. 1.9) as in the case of exact data-constraints [7,8], merely the determination of the Lagrange parameters is modified due to the presence of noise to

$$\begin{aligned} C_{ij} + \alpha \sigma_{ij}^2 &= \int \rho(\eta) \eta_i \eta_j d\eta \\ &= \frac{1}{2}(\lambda^{-1})_{ij} \quad . \end{aligned} \quad (1.17)$$

This relation agrees with Eq. 1.11 for $\sigma_{ij} = 0$. But in practice the errors of the covariance matrix are considerable. Particular if only few QMC data are available the inclusion of the errors of the covariance matrix are essential to obtain an unbiased estimator for the dynamic quantities: The covariance matrix C_{ij} is determined from QMC data D_i^ν via $C_{ij} = \frac{1}{N-1} \sum_{\nu=1}^N \Delta D_i^\nu \Delta D_j^\nu$, where ν represents the independent measurements (bins) of D_i , as discussed below. It is obvious that the rank of C is less or equal to the number of bins (N). Hence, if the number of bins is less than the dimension of the covariance matrix, the inverse of C , entering Eq. 1.12, does not exist, and the regularization term in 1.17 is essential (no matter how small σ_{ij} is) to determine $\rho(\eta)$.

Hence the *Likelihood* $P(D|ACH)$ remains a normal distribution, merely the covariance is not the QMC error-covariance, it rather has to be determined via Eq. 1.17. The regularization parameter α entering Eq. 1.17 can be determined either selfconsistently upon maximizing the marginal *posterior* $P(\alpha|C\sigma H)$ [2,3], or via the historic condition $\chi^2 = N$. We employ the historic approach since we know that the number of good degrees of freedom is small and both stopping criteria will yield essentially the same result [3].

II. APPLICATION TO THE SPECTRAL PROPERTIES OF STRONGLY CORRELATED ELECTRONS

As a typical and topical problem we study the dynamic properties of the Hubbard model which is presently the subject of intense analytical and numerical studies. The detailed understanding of the dynamic properties of strongly correlated electrons is essential for the theoretical description of the high temperature superconductors. The Hubbard model reads

$$H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + h.c.) + U \sum_i n_{i\uparrow} n_{i\downarrow} \quad (2.1)$$

with the hopping matrix element t between two adjacent sites. $c_{i,\sigma}^{(\dagger)}$ destroys (creates) an electron of spin σ on site i , $\langle i, j \rangle$ denotes nearest neighbors, U is a Coulomb repulsion for two electrons of opposite spin on the same site and $n_{i,\sigma} = c_{i,\sigma}^\dagger c_{i,\sigma}$.

Unfortunately, dynamic properties cannot be measured directly by QMC simulations. Dynamical information is provided by Matsubara Greens functions $D_l = - \langle T_\tau P(\tau_l) Q(0) \rangle$ for discrete τ_l -values on the imaginary time axis, where $l = n * \beta / L$, $n = 0, \dots, L$ and L is the number of time slices. P and Q are operators which define the correlation function. Here we will consider the one-particle properties of strongly correlated fermions, and the operators are therefore $P = c$ and $Q = c^\dagger$, respectively. In order to determine the spectral density $A(\omega)$ for real frequencies ω the spectral theorem is applied:

$$D_l = - \int A(\omega) \frac{e^{-\tau_l \omega}}{1 + e^{-\beta \omega}} d\omega \quad (2.2)$$

which is, as already stated above, an inverse Laplace transformation problem and pathologically ill-posed.

Further information on the spectrum is provided by making use of the lowest order moments of $A(\omega)$,

$$\mu_m = \int \omega^m A(\omega) d\omega, \quad (2.3)$$

which are given by commutation relations, $\mu_m = \langle [c, H]_m, c^\dagger \rangle$, and are of simple shape for $m=1,2$ [13].

III. RESULTS

In order to compare the QMC/ME-data with exact results, we consider a chain of $N = 12$ sites, which is still accessible by exact diagonalization (ED) techniques. The QMC-simulations were done for an inverse temperature of $\beta t = 20$ ($T = 0.05t$), where ground state behavior is achieved for this system size and a comparison with the $T = 0$ -ED results is possible. After obtaining thermal equilibrium, up to 640000 sweeps through the space-time Ising-fields were performed.

Data from consecutive measurements are highly correlated, even for the same statistical variable D_l . A study of the skewness (third moment) and the kurtosis (fourth moment) of the data showed, that to get Gaussian behavior at least 200 measurements, each separated by 4 sweeps, have to be accumulated to form one bin. Then the results of a certain number of bins are used for the inversion process.

But binning the data does not suffice to get rid of all the correlations. Still one has to consider the correlations in imaginary time τ (i.e. between D_l and $D_{l'}$). In Fig. 1 the QMC-result for one single bin is compared to the final shape of the Greens function for $N^{bin} = 800$. Instead of being distributed 'at random' around the average the data for this bin are systematically lower than the average for $\tau < \beta/2$ and systematically higher for $\tau > \beta/2$. These correlations may be reduced by forming larger and larger bins (i.e. using more and more computation time). But it is the aim of this paper to show that this is not a sensible thing to do and that one can do better by taking into account the correlations [14] and particularly by accounting for the statistical errors of the covariance matrix.

In the following we will discuss the results of the MaxEnt-procedure considering three cases: (1) Neglecting any information of the covariance matrix, (2) using the covariance matrix only, and (3) taking into account both the covariance and its errors. To determine the dependence on the number of bins and to give an quantitative argument for the amount of the computational effort, which has to be taken, we show spectra resulting from QMC-data for 200, 400 and 800 bins (160000, 320000 and 640000 sweeps, respectively). The

number of time-slices, which corresponds to the dimension of the covariance matrix, is 160 in the present study.

Starting with 200 bins, which is slightly larger than the number of time slices, one can see (first column in Fig. 2) that neither the MaxEnt reconstruction of the plain data (Fig. 2a) nor the additional use of the covariance matrix (Fig. 2d) gives a reliable result. In both spectra the structures are too pronounced and at the wrong position. It appears that the results are generally better if the covariance matrix is not included, since the additional information is treated as exact data-constraints although it suffers from pronounced statistical noise. If, however, the statistical errors of the covariance matrix are taken into account (Fig. 2g) the result reproduces the ED-result very well. There is a small overestimation of the spectral weight at $\omega \approx -3$ only.

Increasing the number of bins to 400 gives still an overfitted result for the first case (Fig. 2b). Taking the covariance matrix into account (Fig. 2e) shows a slightly improved spectrum for $\omega > 0$, but for $\omega \approx -2$ the spectral weight is suppressed completely. Again the best spectrum is obtained if the errors of the covariance matrix are properly accounted for (Fig. 2h). The maximum at $\omega \approx -3$ is damped to the correct shape and the agreement with the ED-result is nearly perfect now.

Eventually for the large number of 800 bins all three spectra show satisfactory results (third column of Fig. 2). Only in the first case (Fig. 2c) the ME-curve decreases still too fast for $\omega > 5$ leading to a wrong width.

The convergency of the various approaches is reasonable, since with increasing number of bins, the correlation of the QMC-errors for different imaginary times vanishes and the covariance matrix becomes diagonal and the covariance of the errors can be ignored. At the same time, the errors of the covariance matrix decrease and assuming exact data-constraints becomes also exact.

Further investigations of the effect of the error of the covariance matrix revealed that the procedure can be simplified by assuming a sufficiently large constant relative error (in our case 20%). The results show no significant deviation from the results obtained by taking

the correctly generated errors.

IV. CONCLUSION

We have shown that the imaginary time data, obtained by standard QMC simulations, suffer from strongly correlated statistical errors if only a small sample is used. It appears reasonable to include this additional knowledge (covariance matrix). But it has been generally observed, the QMC data of the covariance matrix are useless if the sample size is small and they are needless if the sample size is large. And it seemed generally advantageous to ignore the off-diagonal elements of the covariance matrix altogether. At any rate, this approach demands extremely long QMC-runs (more than half a million sweeps was not even sufficient in our case) and gets increasingly impossible for larger system sizes (the computation time for one sweep scales with $\sim N^3$). We have shown that only if the covariance matrix and its errors are treated consistently in the Bayesian frame reliable results can be obtained regardless of the number bins.

V. ACKNOWLEDGMENTS

We would like to thank R. Silver and S.R. White for stimulating discussions and we are grateful to the Bavarian "FORSUPRA" program on high T_c research for financial support. The calculations were performed at HLRZ Jülich and at LRZ München.

REFERENCES

- [1] A. O'Hagan, "Kendall's advanced theory of statistics", 2b, "Bayesian Inference", Halsted Press ISBN 047023388I 8, (1994).
- [2] J. Skilling, in *Maximum Entropy and Bayesian Methods*, ed. J. Skilling (Kluwer Academic Publishers, Dordrecht, 1989), p.45; J. Skilling, in *Maximum Entropy and Bayesian Methods* ed. P. F. Fougère, (Kluwer, Academic Publishers, 1990), p.341.
- [3] S.F. Gull, in *Maximum Entropy and Bayesian Methods* ed. J. Skilling, (Kluwer, Academic Publishers, 1989), p.53.
- [4] *Maximum Entropy in Action*, ed. B. Buck and V. A. Macaulay, (Oxford Science Publications, Oxford, 1990).
- [5] R.N. Silver, D.S. Sivia, and J.E. Gubernatis, Phys. Rev. B**41**, 2380 (1990), and references therein.
- [6] W. von der Linden, M. Donath, and V. Dose, Phys. Rev. Lett.**71**,899 (1993).
- [7] R.N. Silver and H.F. Martz, in *Maximum Entropy and Bayesian Methods* ed. G. Heidbreder, (Kluwer, Academic Publishers, 1993), to be published.
- [8] Wolfgang von der Linden, Appl.Phys. **A60**, 155 (1994)
- [9] R.N. Silver, et al., Phys. Rev. Lett. **65**, 496 (1990).
- [10] J. Skilling, in *Maximum Entropy in Action*, ed. B. Buck and V. A. Macaulay, (Oxford Science Publications, Oxford, 1990), p. 19.
- [11] *Entropy Optimization Principles with Applications*, eds. J.N. Kapur and H.K. Kesavan, (Academic Press, Inc., San Diego, 1992).
- [12] J.E. Gubernatis, et al., Phys. Rev. B**44**, 6011 (1991).
- [13] S.R. White, Phys. Rev. B**44**, 4670 (1991).

[14] In this section we followed suggestions of a talk manuscript by Mark Jarrell, University of Cincinnati.

FIGURES

FIG. 1. The data from one bin (points) compared to the final average over 800 bins (line).

FIG. 2. Comparison of the MaxEnt-spectra (thick line) with the ED-result (thin line) for different parameters: first row (a,b,c): without use of the covariance (first case, see text), second row (d,e,f): with covariance (second case), third row (g,h,i): with the error of the covariance matrix (third case); in dependence of the number of bins: first column (a,d,g): 200 bins, second column (b,e,h): 400 bins, third column (c,f,i): 800 bins.




Fig. 1




Fig. 2